

PDE and Boundary-Value Problems

Winter Term 2014/2015

Lecture 20

Saarland University

5. Februar 2015

Purpose of Lesson

- To introduce the idea of explicit finite-difference methods and show how they can be used to solve hyperbolic and parabolic problems.
- To show how time-dependent problems can be solved by another finite-difference scheme known as **implicit methods**.

- We can solve elliptic BVPs (steady-state problems) where the PDE was satisfied in a given region of space, and the solution (or its derivative) was specified on the boundary.
- In those types of problems, we find the approximate solution at the **interior grid points** by solving a system of algebraic equations. In other words, the solution at all the interior grid points was found **simultaneously**.

An Explicit Finite-Difference Method

- Now we will show how **time-dependent problems** can be solved by finite-difference approximations.
- The idea is that if we are given the solution when time is **zero**, we can then find the solution for $t = \Delta t, 2\Delta t, 3\Delta t, \dots$ by means of a **marching process**.
- Replacing both the **space** and **time** derivatives by their finite-difference approximations, we can then solve for the solution $u_{i,j}$ in the difference equation **explicitly** in terms of the solution at earlier values of time.
- This process is called an **explicit-type marching process**, since we find the solution at a **single** value of time in terms of the solution at earlier values of time.

The Explicit Method for Parabolic Equations

- To show how the explicit finite-difference method works, we consider a representative problem from heat flow.
- Heat flows along a rod initially at temperature zero, where the left end of the rod is fixed at temperature one, and the right-hand side experiences a heat loss (or gain) proportional to the difference between the temperature at that end and an outside temperature that is given by $g(t)$.

The Explicit Method for Parabolic Equations (cont.)

Problem 20-1

To find a function $u(x, t)$ that satisfies

$$\text{PDE: } u_t = u_{xx}, \quad 0 < x < 1, \quad 0 < t < \infty$$

$$\text{BCs: } \begin{cases} u(0, t) = 1 \\ u_x(1, t) = -[u(1, t) - g(t)] \end{cases} \quad 0 < t < \infty$$


$$\text{IC: } u(x, 0) = 0 \quad 0 \leq x \leq 1$$

To solve problem 20-1 by finite differences, we start by drawing the usual rectangular grid system with grid points

$$x_j = jh \quad j = 0, 1, 2, \dots, n$$

$$t_i = ik \quad i = 0, 1, 2, \dots, m$$

Grid system for a heat-flow problem



- Note that on the figure of the grid system, the $u_{i,j}$ on the **left** and **bottom** are given BCs and ICs, and our job is to find the other $u_{i,j}$'s.
- To do this, we begin by replacing the partial derivatives u_t and u_{xx} in the heat equation with their approximations

$$u_t = \frac{1}{k} [u(x, t + k) - u(x, t)] = \frac{1}{k} (u_{i+1,j} - u_{i,j})$$

$$\begin{aligned} u_{xx} &= \frac{1}{h^2} [u(x + h, t) - 2u(x, t) + u(x - h, t)] \\ &= \frac{1}{h^2} (u_{i,j+1} - 2u_{i,j} + u_{i,j-1}) \end{aligned}$$

- By substituting these approximations into $u_t = u_{xx}$ and solving for the solution at the largest value of time, we have

$$u_{i+1,j} = u_{i,j} + \frac{k}{h^2} [u_{i,j+1} - 2u_{i,j} + u_{i,j-1}] \quad (20.1)$$

Remark

(20.1) is the formula we are looking for, since it gives us the solution at one value of time in terms of the solution at earlier values of time.

- We are almost ready to begin the computations for problem 20-1. First, we must approximate the derivatives in the right-hand BC

$$u_x(1, t) = - [u(1, t) - g(t)]$$

by

$$\frac{1}{h} [u_{i,n} - u_{i,n-1}] = - [u_{i,n} - g_i], \quad (20.2)$$

where $g_i = g(ik)$ is given.

- Note that in (20.2) we have replaced $u_x(1, t)$ by the **backward-difference approximation**, since the forward-difference approximation would require knowing values of $u_{i,j}$ outside the domain.
- Solving (20.2) for $u_{i,n}$ gives us

$$u_{i,n} = \frac{u_{i,n-1} + hg_i}{1 + h}. \quad (20.3)$$

Algorithm for the Explicit Method

1. Find the solution at the grid points for $t = \Delta t$ by using the explicit formula

$$u_{2,j} = u_{1,j} + \frac{k}{h^2} [u_{1,j+1} - 2u_{1,j} + u_{1,j-1}] \quad j = 2, 3, \dots, n-1$$


2. Find $u_{2,n}$ from formula (20.3)

$$u_{2,n} = \frac{u_{2,n-1} + hg_2}{1 + h}.$$

Remark

- Steps 1 and 2 find the solution for $t = \Delta t$.
- To find the solution for $t = 2\Delta t$ repeat steps 1 and 2, moving up one more row (increase i by 1) and using the values of $u_{i,j}$ just computed.
- For $t = 3\Delta t, 4\Delta t, \dots$ keep repeating the same process.

On the flow diagram on the next page we explain in a precise manner how the computations should be carried out.



Remarks

- There is a serious deficiency in the explicit method, for if the step size in t is large compared to the step size in x , then machine roundoff error can grow until it ruins the accuracy of the solution.
- The relative size of these steps depends on the particular equation and the BCs, but, generally, the step size in t should be much smaller than the step size in x . We must have $k/h^2 \leq 0.5$ in order this method to work.
- A general rule of thumb is that as the step sizes Δt and Δx are made smaller, the **truncation error** of approximating partial derivatives by finite differences decreases. However, the smaller these grid sizes, the more computations necessary, and, hence, the **roundoff error**, as a result of rounding off our computations, will increase.

The Explicit Method for Hyperbolic Equation

Problem 20-2

To find a function $u(x, t)$ that satisfies

$$\text{PDE: } u_{tt} = u_{xx}, \quad 0 < x < 1, \quad 0 < t < \infty$$

$$\text{BCs: } \begin{cases} u(0, t) = g_1(t) \\ u(1, t) = g_2(t) \end{cases} \quad 0 < t < \infty$$

$$\text{ICs: } \begin{cases} u(x, 0) = \phi(x) \\ u_t(x, 0) = \psi(x) \end{cases} \quad 0 \leq x \leq 1$$

- Problem 20-2 can also be solved by the explicit finite-difference method. Here, we can approximate the derivatives u_{tt} and u_{xx} by

$$u_{tt} \cong \frac{1}{k^2} [u(x, t+k) - 2u(x, t) + u(x, t-k)]$$
$$u_{xx} \cong \frac{1}{h^2} [u(x+h, t) - 2u(x, t) + u(x-h, t)]$$

and the derivative $u_t(x, 0)$ in the IC by

$$u_t(x, 0) \cong \frac{1}{k} [u(x, k) - u(x, 0)] = \frac{1}{k} [u(x, k) - \phi(x)].$$

- Solving for $u(x, t + k)$ explicitly in terms of the solution at earlier values of time gives

$$u(x, t + k) = 2u(x, t) - u(x, t - k) + \left(\frac{k}{h}\right)^2 [u(x + h, t) - 2u(x, t) + u(x - h, t)] \quad (20.4)$$

- From (20.4) it is clear that we must already know the solution at **two** previous time steps, and, hence, we must use the initial-velocity condition

$$\frac{1}{k} [u(x, k) - \phi(x)] = \psi(x)$$

to get us started. Solving for $u(x, k)$ gives $u(x, k) = \phi(x) + k\psi(x)$, and, thus, we can find the solution for $t = \Delta t$.

An Implicit Finite-Difference Method (Crank-Nicolson Method)

- In implicit method, we again replace the partial derivatives in the problem by their finite-difference approximations, but unlike explicit methods (where we solved for $u_{i+1,j}$ explicitly in terms of earlier values), in implicit methods, we solve a **system of equations** in order to find the solution at the largest value of time.
- In other words, for each new value of time we solve a system of algebraic equations to find **all** the values.
- It should be mentioned that implicit methods allow us to take larger steps by doing more work per step.

The Heat-Flow Problem Solved by an Implicit Method


Problem 20-3

To find a function $u(x, t)$ that satisfies

$$\text{PDE: } u_t = u_{xx}, \quad 0 < x < 1, \quad 0 < t < \infty$$

$$\text{BCs: } \begin{cases} u(0, t) = 0 \\ u(1, t) = 0 \end{cases} \quad 0 < t < \infty$$

$$\text{IC: } u(x, 0) = 1 \quad 0 \leq x \leq 1$$

Grid system for implicit scheme ($\Delta x=0.2$)

- We replace the partial derivatives u_t and u_{xx} by the following approximations:

$$u_t(x, t) = \frac{1}{k} [u(x, t + k) - u(x, t)]$$

$$u_{xx}(x, t) = \frac{\lambda}{h^2} [u(x + h, t + k) - 2u(x, t + k) + u(x - h, t + k)] \\ + \frac{(1 - \lambda)}{h^2} [u(x + h, t) - 2u(x, t) + u(x - h, t)],$$

where λ is a chosen number in the interval $[0, 1]$.

- Note that our approximation for u_{xx} is a **weighted average** of the central-difference approximation to the derivative u_{xx} at time values t and $t + k$.

Remarks

- In the special case when $\lambda = 0.5$, it is just the ordinary average of these two central differences.
- If $\lambda = 0.75$, our approximation puts weights of 0.75 and 0.25 on each of the two terms.
- If $\lambda = 0$, it is usual **explicit** finite-difference method.

If we now substitute the approximations for u_t and u_{xx} into our problem, we get the new **finite-difference problem**

Problem 20-3a

$$\frac{1}{k} (u_{i+1,j} - u_{i,j}) = \frac{\lambda}{h^2} (u_{i+1,j+1} - 2u_{i+1,j} + u_{i+1,j-1}) + \frac{(1-\lambda)}{h^2} (u_{i,j+1} - 2u_{i,j} + u_{i,j-1})$$

$$\text{BCs: } \begin{cases} u_{i,1} = 0 \\ u_{i,n} = 0 \end{cases}, \quad i = 1, 2, \dots, m$$

$$\text{IC: } u_{1,j} = 1, \quad j = 2, \dots, n-1$$

- If we rewrite the difference equation in problem 20-3a, putting the $u_{i,j}$'s with the largest time subscript (i -subscript) on the left-hand side of the equation, we arrive at

$$\begin{aligned}
 -\lambda r u_{i+1,j+1} + (1 + 2r\lambda)u_{i+1,j} - \lambda r u_{i+1,j-1} \\
 = r(1 - \lambda)u_{i,j+1} + [1 - 2r(1 - \lambda)] u_{i,j} \\
 + r(1 - \lambda)u_{i,j-1},
 \end{aligned} \tag{20.5}$$

where we have set $r = k/h^2$ for convenience.

- Note that for a **fixed subscript i** and for j going from 2 to $n - 1$, this is a system of $n - 2$ equations in the $n - 2$ unknowns $u_{i+1,2}$, $u_{i+1,3}$, $u_{i+1,4}$, \dots , $u_{i+1,n-1}$ [which are the interior grid points at $t = (i + 1)\Delta t$].
- To help show exactly how $u_{i,j}$'s are involved into (20.5), we write it in the symbolic or molecular form (see next page)

The molecule form of the implicit formula

