

Calculus of Variations

Summer Term 2014

Lecture 11

6. Juni 2014

Purpose of Lesson:

- To introduce the notion of a broken extremal
- To discuss the properties of broken extremals

§7. Broken extremals

- Until now we mostly studied the extremal curves with at least two well-defined derivatives.
- Obviously this is not always true.
- **Broken extremals** are continuous extremals for which the gradient has a discontinuity at one or more points.
- If a variational problem has a smooth extremal (That therefore satisfies the Euler-Lagrange equations), this will be better than a broken one.
- But some problems don't admit smooth extremals.

Example 11.1

Find $y(x)$ to minimize

$$J[y] = \int_{-1}^1 y^2(1 - y')^2 dx$$

subject to $y(-1) = 0$ and $y(1) = 1$.

Example 11.1 (cont.)

- There is no explicit x dependence inside the integral, so we can find

$$f - y'f_{y'} = c_1 = \text{const}$$

$$y^2(1 - y')^2 + 2y'y^2(1 - y') = c_1$$

$$y^2(1 - y') [1 + y'] = c_1$$

$$y^2 [1 - y'^2] = c_1$$

- If $c_1 = 0$ we get the singular solutions

$$y = 0 \quad \text{or} \quad y = \pm x + B.$$

Neither of these satisfies both end-points conditions $y(-1) = 0$ and $y(1) = 1$, so $c_1 \neq 0$ (we think).

Example 11.1 (cont.)

- Given $c_1 \neq 0$

$$y^2 [1 - y'^2] = c_1$$

$$y'^2 = \frac{y^2 - c_1}{y^2}$$

$$\frac{dy}{dx} = \pm \frac{1}{y} \sqrt{y^2 - c_1}$$

$$dx = \pm \frac{y}{\sqrt{y^2 - c_1}} dy$$

$$x = \pm \sqrt{y^2 - c_1} + c_2$$

$$(x - c_2)^2 = y^2 - c_1$$

- The solution is a **rectangular hyperbola**.

Example 11.1 (cont.)

- Using the end-points conditions we find c_1 and c_2 from

$$(x - c_2)^2 = y^2 - c_1.$$

$$y(-1) = 0 \quad \Rightarrow \quad (-1 - c_2)^2 = -c_1$$

$$y(1) = 1 \quad \Rightarrow \quad (1 - c_2)^2 = 1 - c_1$$

- Addition of these two equations gives


$$(1 - c_2)^2 = 1 + (1 + c_2)^2$$

which has solution $c_2 = -1/4$, and so $c_1 = -9/16$

$$y^2 = (x + 1/4)^2 - 9/16.$$

Example 11.1 (cont.)


- The end-points are on opposite branches of the hyperbola!



- There is **NO** smooth extremal curve that connects $(-1, 0)$ and $(1, 1)$.

- Sometimes there is no **smooth** extremal.
- We must seek a **broken extremal**.
- Still want a continuous extremal.
- What should we do?
 - Previous smoothness results suggest that we should use a smooth extremal when we can, and so we will try to minimize the number of **corners**.
 - We'll start by looking for curves with one corner.
 - But can we apply the Euler-Lagrange equations?

- If we have an extremal like this, can we use the Euler-Lagrange equations?




Theorem 11.1

If the smooth curve $y(x)$ gives an extremal of a functional $J[y]$ over the class of all admissible curves in some ε neighborhood of y , then $y(x)$ also gives an extremal of a functional $J[y]$ over the class of all **piecewise smooth curves** in the same neighborhood.

Meaning:

We can extend our results to piecewise smooth curves (where a smooth result exists), not just curves with 2 continuous derivatives.




- The theorem assumes that there exists a smooth extremal (in this case a minimum for the purpose of illustration) y . Then for any other smooth curve $\hat{y} \in B_\varepsilon(y)$ we know $J[\hat{y}] > J[y]$.
- Assume for the moment that for a piecewise smooth function $\tilde{y} \in B_\varepsilon(y)$ we have $J[\tilde{y}] < J[y]$. We can approximate \tilde{y} by a smooth curve $\hat{y}_\delta \in B_\varepsilon(y)$ by rounding off the edges of the discontinuity.
- Given that we can approximate the curve \tilde{y} arbitrarily closely by a smooth curve \hat{y}_δ , for which we already know $J[\hat{y}_\delta] > J[y]$. We get a contradiction with $J[\tilde{y}] < J[y]$, and so no such alternative extremal can exist.


- Break the functional into two parts:

$$J[y] = J_1[y] + J_2[y] = \int_{x_0}^{x^*} F(x, y_1, y_1') dx + \int_{x^*}^{x_1} F(x, y_2, y_2') dx$$

- We require y to have two continuous derivatives everywhere except at x^* , and $y_1(x^*) = y_2(x^*)$.



Possible perturbations:



The location of the "corner" can also be perturbed

- We get the first component of the first variation by considering a problem with only one fixed end-point, and allowing x^* to vary, so that

$$\begin{aligned}
 0 &= \left. \frac{d\phi_1(\varepsilon)}{d\varepsilon} \right|_{\varepsilon=0} = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \int_{x_0}^{\hat{x}^*} F(x, y_1 + \varepsilon\eta, y_1' + \varepsilon\eta') dx \\
 &= \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \int_{x_0}^{x^* + \varepsilon X} F(x, y_1 + \varepsilon\eta, y_1' + \varepsilon\eta') dx \\
 &= \left. XF(x, y_1, y_1') \right|_{x=x^*} + \left. F_{y_1'}\eta \right|_{x=x^*} + \int_{x_0}^{x^*} \left(F_{y_1} - \frac{d}{dx} F_{y_1'} \right) dx
 \end{aligned}$$

- The perturbed point (\hat{x}^*, \hat{y}^*) and perturbed function η must satisfy certain conditions to be compatible.
- Remember that

$$\hat{x}^* = x^* + \varepsilon X$$

$$\hat{y}^* = y^* + \varepsilon Y$$

- Notice that

$$\hat{y}^* = y(x^* + \varepsilon X) + \varepsilon \eta(x^* + \varepsilon X).$$

- From Taylor's theorem, for small ε

$$y(x^* + \varepsilon X) = y(x^*) + \varepsilon X y'(x^*) + O(\varepsilon^2)$$

$$= y^* + \varepsilon X y'(x^*) + O(\varepsilon^2)$$

$$\varepsilon \eta(x^* + \varepsilon X) = \varepsilon \eta(x^*) + O(\varepsilon^2)$$

- So

$$y^* + \varepsilon Y = y^* + \varepsilon Xy'(x^*) + \varepsilon \eta(x^*) + O(\varepsilon^2)$$

$$\varepsilon Y = \varepsilon Xy'(x^*) + \varepsilon \eta(x^*) + O(\varepsilon^2)$$

$$\eta(x^*) = Y - Xy'(x^*) + O(\varepsilon)$$

- Thus, we have

$$\boxed{\eta(x^*) = Y - Xy'(x^*) + O(\varepsilon)} \quad (11.1)$$

- Substituting the compatibility constraint (11.1) into the our first variation we get

$$\begin{aligned}
 0 &= \left[XF + F_{y_1'} \eta \right]_{x=x^*} + \int_{x_0}^{x^*} \left(F_{y_1} - \frac{d}{dx} F_{y_1'} \right) \eta dx \\
 &= XF|_{x=x^*} + [Y - Xy_1'(x^*)] F_{y_1'}|_{x=x^*} + \int_{x_0}^{x^*} \left(F_{y_1} - \frac{d}{dx} F_{y_1'} \right) \eta dx \\
 &= X \left[F - y_1' F_{y_1'} \right]_{x=x^*} + Y F_{y_1'}|_{x=x^*} + \int_{x_0}^{x^*} \left(F_{y_1} - \frac{d}{dx} F_{y_1'} \right) \eta dx
 \end{aligned}$$

- So, we get an integral term which results in the E-L equation, plus the additional constraint

$$X \left[F - y_1' F_{y_1'} \right]_{x=x^*} + Y F_{y_1'}|_{x=x^*} = 0 \quad (11.2)$$

- Note that, for the second component of the First Variation we get a similar extra term, e.g.

$$-X \left[F - y_2' F_{y_2'} \right]_{x=x^*} - Y F_{y_2'} \Big|_{x=x^*} = 0. \quad (11.3)$$

- The sign is reversed because it corresponds to the x_0 term (as opposed to the x_1 term for δJ_1).
- The combined First Variation (minus the terms that result from the Euler-lagrange equation which must be zero) is

$$X \left[F - y_1' F_{y_1'} \right]_{x=x^*} + Y F_{y_1'} \Big|_{x=x^*} - X \left[F - y_2' F_{y_2'} \right]_{x=x^*} - Y F_{y_2'} \Big|_{x=x^*} = 0.$$

- We rearrange to give

$$0 = X \left\{ \left[F(x, y_1, y'_1) - y'_1 F_{y'_1} \right] - \left[F(x, y_2, y'_2) - y'_2 F_{y'_2} \right] \right\}_{x=x^*} \\ + Y \left\{ F_{y'_1} - F_{y'_2} \right\}_{x=x^*} .$$

- Note that the point of discontinuity may vary freely, so we may independently vary X and Y or set one or both to zero. Hence, we can separate the condition to get two conditions

$$\left[F(x, y_1, y'_1) - y'_1 F_{y'_1} - F(x, y_2, y'_2) + y'_2 F_{y'_2} \right]_{x=x^*} = 0 \\ \left\{ F_{y'_1} - F_{y'_2} \right\}_{x=x^*} = 0$$

- We can write the conditions as

$$\begin{aligned} \left[F(x, y_1, y_1') - y_1' F_{y_1'} \right]_{x=x^*} &= \left[F(x, y_2, y_2') - y_2' F_{y_2'} \right]_{x=x^*} \\ F_{y_1'} \Big|_{x=x^*} &= F_{y_2'} \Big|_{x=x^*} \end{aligned}$$

Called the **Weierstrass-Erdmann Corner Conditions**.

- Rather than separating y into y_1 and y_2 we may write the corner conditions in terms of limits from the left and right, e.g.

$$\begin{aligned} \left[F - y' F_{y'} \right]_{x=x^{*-}} &= \left[F - y' F_{y'} \right]_{x=x^{*+}} \\ F_{y'} \Big|_{x=x^{*-}} &= F_{y'} \Big|_{x=x^{*+}} \end{aligned}$$

So the broken extremal solution must satisfy

- The Euler-Lagrange equations
- The Weierstrass-Erdmann Corner Conditions

$$\begin{aligned} [F - y'F_{y'}]_{x=x^{*-}} &= [F - y'F_{y'}]_{x=x^{*+}} \\ F_{y'} \Big|_{x=x^{*-}} &= F_{y'} \Big|_{x=x^{*+}} \end{aligned}$$

must hold at any "corner".

Example 11.1 (cont. II)

Find $y(x)$ to minimize

$$J[y] = \int_{-1}^1 y^2(1 - y')^2 dx$$

subject to $y(-1) = 0$ and $y(1) = 1$.

Example 11.1 (cont. II)

- In the example considered

$$F - y' F_{y'} = y^2 (1 - y'^2)$$
$$F_{y'} = -2y^2 (1 - y')$$

- Remember that $y = 0$ and $y = x + A$ are valid solutions to the Euler-Lagrange equations, and that for both of these solutions

$$F_{y'} = F - y' F_{y'} = 0,$$

so we can put a "corner" where needed.

Example 11.1 (cont. II)

- The solution must also satisfy the end-point conditions, so $y(-1) = 0$ and $y(1) = 1$, and therefore, as valid solution has $x^* = 0$ and

$$y_1 = 0 \quad \text{for} \quad x \in [-1, x^*]$$

$$y_2 = x \quad \text{for} \quad x \in [x^*, 1]$$