
Stochastics II

10. Tutorial

Let $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \in [0, \infty)}, P)$ be the underlying filtered probability space for the whole assignment.

Exercise 1 (3 Points) Let X be an \mathbb{F} -adapted and integrable stochastic process with right-continuous paths. Show that X is a submartingale, if and only if the equality $E[X_\sigma] \leq E[X_\tau]$ holds for every bounded stopping times σ and τ with $\sigma \leq \tau$.

Exercise 2 (3 Points) Show that if the stochastic process W is a Brownian motion with respect to the filtration \mathbb{F} , then it is also a Brownian motion with respect to \mathcal{F}^+ .

Exercise 3 (7 Points) Let $X = (X_t)_{t \in [0, \infty)}$ be a stochastic process with continuous paths, starting at $X_0 = 0$.

(i) Show that X is a Brownian motion with respect to \mathbb{F} , if and only if X is \mathbb{F} -adapted and

$$E[e^{iu(X_t - X_s)} | \mathcal{F}_s] = e^{-\frac{1}{2}u^2(t-s)}$$

for every $0 \leq s < t$ and $u \in \mathbb{R}$.

(ii) Show that X is a Brownian motion with respect to \mathbb{F}^X , if and only if X is a Gaussian process with $E[X_t] = 0$ and $Cov(X_t, X_s) = t \wedge s$, for $s, t \in [0, \infty)$.

Exercise 4 (4 Points) Let $X = (X_t)_{t \in [0, \infty)}$ be a Brownian motion with respect to \mathbb{F} and $a, b > 0$. We take a look at the Gaussian process $Y = (Y_t)_{t \in [0, \infty)}$, given by

$$\forall t \in [0, \infty) : Y_t = \sqrt{a}e^{-bt}X_{e^{2bt}}.$$

Calculate $E[Y_t]$ and $Cov(Y_t, Y_s)$ for $s, t \in [0, \infty)$.