

## Stochastics II

### 12. Tutorial

**Definition** A function  $\varrho : \mathcal{T} \times \mathcal{T} \rightarrow \mathbb{R}$  is called positive semi-definite, if

$$\forall k \in \mathbb{N} \forall t_1, \dots, t_k \in \mathcal{T} \forall \lambda_1, \dots, \lambda_k \in \mathbb{R} : \sum_{i=1}^k \sum_{j=1}^k \lambda_i \lambda_j \varrho(t_i, t_j) \geq 0.$$

**Exercise 1** (5 Points) Let  $m : \mathcal{T} \rightarrow \mathbb{R}$  be arbitrary and let  $\varrho : \mathcal{T} \times \mathcal{T} \rightarrow \mathbb{R}$  be symmetric and positive semi-definite. Show that there exists a probability space  $(\Omega, \mathcal{F}, P)$  and a Gaussian process  $X = (X_t)_{t \in \mathcal{T}}$  on  $(\Omega, \mathcal{F}, P)$  such that  $E[X_t] = m(t)$  and  $\text{Cov}(X_s, X_t) = \varrho(s, t)$  for all  $s, t \in \mathcal{T}$ .

**Exercise 2** (1+3+2+4 Points) Let  $T \in \mathbb{R}$  and  $(X_t)_{t \in [0, T]}$  be the stochastic process defined by

$$X_t := \frac{T-t}{T} W_{\frac{tT}{T-t}}$$

for every  $t \in \mathbb{R}_+$ , where  $(W_t)_{t \in [0, \infty)}$  is a Brownian motion.

- (i) Show that  $X_T := \lim_{t \nearrow T} X_t$   $P$ -a.s. exists and derive the distribution of  $X_T$ .
- (ii) Show that  $(X_t)_{t \in [0, T]}$  is a Gaussian process and calculate  $E[X_t]$  and  $\text{Cov}(X_s, X_t)$  for  $s, t \in [0, T]$ .
- (iii) Show that the random variables

$$Z_i = \frac{X_{t_i}}{T - t_i} - \frac{X_{t_{i-1}}}{T - t_{i-1}}$$

are independent for  $k \in \mathbb{N}$ ,  $t_1, \dots, t_k \in [0, T)$ .

- (iv) Show that

$$P \left( \bigcap_{i=1}^k \{X_{t_i} \leq x_i\} \right) = \lim_{\varepsilon \searrow 0} P \left( \bigcap_{i=1}^k \{W_{t_i} \leq x_i\} \mid \{W_T \in (-\varepsilon, \varepsilon)\} \right)$$

for  $k \in \mathbb{N}$ ,  $t_1, \dots, t_k \in [0, T)$  and  $x_1, \dots, x_k \in \mathbb{R}$ .

**Exercise 3** (4+1Points)

- (i) Show that

$$\frac{x}{1+x^2} e^{-\frac{x^2}{2}} \leq \int_x^\infty e^{-\frac{t^2}{2}} dt \leq \frac{1}{x} e^{-\frac{x^2}{2}}$$

for every  $x > 0$  and conclude that

$$P(\{\xi > x\}) \geq \frac{1}{\sqrt{2}^3 \sqrt{\pi}} \frac{1}{x} e^{-\frac{x^2}{2}}$$

for every  $\mathcal{N}(0, 1)$ -distributed random variable  $\xi$  and all  $x \geq 1$ .

(ii) We define for  $n \in \mathbb{N}$  and  $q > 1$

$$a_n := \frac{e^{-\log \log(q^n - q^{n-1})}}{\sqrt{\log \log(q^n - q^{n-1})}}.$$

Show that  $\sum_{n \in \mathbb{N}} a_{2n} = \infty$ .