

Universität des Saarlandes



Fachrichtung 6.1 – Mathematik

Preprint Nr. 201

Error estimates for obstacle problems of higher order

Michael Bildhauer and Martin Fuchs

Saarbrücken 2007

Error estimates for obstacle problems of higher order

Michael Bildhauer

Saarland University
Department of Mathematics
P.O. Box 15 11 50
D-66041 Saarbrücken
Germany
bibi@math.uni-sb.de

Martin Fuchs

Saarland University
Department of Mathematics
P.O. Box 15 11 50
D-66041 Saarbrücken
Germany
fuchs@math.uni-sb.de

Edited by
FR 6.1 – Mathematik
Universität des Saarlandes
Postfach 15 11 50
66041 Saarbrücken
Germany

Fax: + 49 681 302 4443
e-Mail: preprint@math.uni-sb.de
WWW: <http://www.math.uni-sb.de/>

AMS Subject Classification: 65 N 15, 65 K 10, 49 J 40, 49 M 29

Keywords: higher order variational inequalities, a posteriori error estimates, duality theory

Abstract

For obstacle problems of higher order involving power growth functionals we prove a posteriori error estimates using methods from duality theory. These estimates can be seen as a reliable measure for the deviation of an approximation from the exact solution being independent of the concrete numerical scheme under consideration.

1 Introduction

In the recent paper [BFR2] we derived a posteriori error estimates for a class of higher order variational inequalities on a planar domain modelling elastic plates with an obstacle subject to a power hardening law. The purpose of the present note is to establish such error estimates for a related type of variational inequalities but under different boundary conditions for which it is also possible to give an estimate of the distance of an arbitrary function satisfying these boundary conditions to the convex set of all admissible comparison functions which respect the obstacle.

To be more precise, let $\Omega \subset \mathbb{R}^2$ denote a bounded smooth domain and introduce the class

$$\mathbb{K} := \{v \in W_p^2(\Omega) \cap \mathring{W}_p^1(\Omega) : v \geq \Psi \text{ on } \Omega\},$$

where $1 < p < \infty$ is fixed and $\mathring{W}_p^1(\Omega)$, $W_p^2(\Omega)$, etc., denote the standard Sobolev spaces, see, e.g. [Ad]. The function Ψ is chosen from $W_p^2(\Omega)$ with the properties $\Psi|_{\partial\Omega} < 0$ and $\Psi(x_0) > 0$ at least for some point $x_0 \in \Omega$. By Sobolev's embedding theorem (compare [Ad]) functions $v \in W_p^2(\Omega)$ are in the space $C^0(\overline{\Omega})$ which immediately shows that \mathbb{K} is non-empty. The variational problem under consideration is

$$(\mathcal{P}) \quad J[v] := \int_{\Omega} \Pi_p(\nabla^2 v) dx + \int_{\Omega} \pi_p(\nabla v) dx \rightarrow \min \text{ in } \mathbb{K},$$

where $\nabla^2 v$ is the matrix of the second generalized partial derivatives of v . Moreover, we have abbreviated

$$\Pi(E) := \frac{1}{p} |E|^p$$

for symmetric (2×2) -matrices E and

$$\pi(\xi) := \frac{1}{p} |\xi|^p$$

Acknowledgement: The authors like to thank S. Repin for many stimulating discussions.

for vectors $\xi \in \mathbb{R}^2$. The reader should note that the variational inequality describing the behaviour of a plate with an obstacle is formulated on the space $\mathring{W}_p^2(\Omega)$ which means that we consider functions with zero trace whose normal derivative also vanishes on $\partial\Omega$. Moreover, for modelling plates it is not necessary to introduce the first order term $\int_{\Omega} \pi_p(\nabla v) dx$ in the functional since we have the coercivity of

$$\mathring{W}_p^2(\Omega) \ni v \mapsto \int_{\Omega} \Pi_p(\nabla^2 v) dx,$$

which is no longer true if $\mathring{W}_p^2(\Omega)$ is replaced by the larger class $W_p^2(\Omega) \cap \mathring{W}_p^1(\Omega)$. Therefore, the weakening of the boundary condition in problem (\mathcal{P}) makes it necessary to include a first order term in the functional J but as the reader can imagine it is not really necessary that $\nabla^2 v$ and ∇v appear with the same power p .

Now, let $u \in \mathbb{K}$ denote the unique solution of problem (\mathcal{P}) . If $v \in \mathbb{K}$ is any comparison function, then we are going to prove an estimate of the form

$$(1.1) \quad \|\nabla^2 u - \nabla^2 v\|_{L^p} + \|\nabla u - \nabla v\|_{L^p} \leq \mathcal{M}(v, \dots),$$

where \mathcal{M} is a non-negative functional depending on v , on the data p , Ω , Ψ and on parameters which are under our disposal. Of course (1.1) is only meaningful provided we can establish the following properties of \mathcal{M} :

- a.) the value of \mathcal{M} is easy to calculate;
- b.) $\mathcal{M}(v, \dots) = 0$ if and only if $v = u$; moreover: $\mathcal{M}(v_k, \dots) \rightarrow 0$ if $v_k \rightarrow u$;
- c.) $\mathcal{M}(v, \dots)$ gives a realistic upper bound for the distance of the approximation v to the exact solution u .

The requirement formulated in c.) means that during the process of deriving (1.1) one should try to avoid overestimation so that (1.1) can be used for a reliable verification of the accuracy of approximative solutions obtained by various numerical methods. We emphasize that the way of how to derive (1.1) is based on purely functional grounds which means that one uses tools from variational calculus such as duality theory which do not refer to any concrete discretization of the problem. Such functional type a posteriori error estimates mainly have been established for a variety of problems by S. Repin, we refer to the monograph [NR] where the interested reader will find further information.

Our paper is organized as follows: in Section 2 we will prove an estimate like (1.1) following [Re] and the modifications of this work outlined in [BFR2]. Here we concentrate on the case $p \geq 2$ since the subquadratic situation requires different techniques, see e.g. [BR] and [BFR1]. In Section 3 we are going to remove the restriction $v \in \mathbb{K}$ from our estimate (1.1) which means that we want to insert arbitrary functions $\tilde{v} \in W_p^2(\Omega) \cap \mathring{W}_p^1(\Omega)$. In order to do so we have to measure the distance of \tilde{v} to the set \mathbb{K} which is possible by using the L^p -theory for elliptic equations.

2 Perturbations of Problem (\mathcal{P})

Let $p \geq 2$, $q := p/(p-1)$, and consider the spaces

$$X := L^p(\Omega; \mathbb{R}^2), \quad Y := L^p(\Omega; \mathbb{R}_{\text{sym}}^{2 \times 2})$$

together with their dual variants

$$X^* = L^q(\Omega; \mathbb{R}^2), \quad Y^* = L^q(\Omega; \mathbb{R}_{\text{sym}}^{2 \times 2})$$

If Π_p^* , π_p^* denote the conjugate functions of Π_p , π_p (see [ET]), then it holds

$$J[w] = \sup_{a^* \in X^*, \tau^* \in Y^*} \int_{\Omega} [a^* \cdot \nabla w + \tau^* : \nabla^2 w - \Pi_p^*(\tau^*) - \pi_p^*(a^*)] dx,$$

and if we introduce the Lagrangian

$$\ell(w, a^*, \tau^*) := \int_{\Omega} [a^* \cdot \nabla w + \tau^* : \nabla^2 w - \Pi_p^*(\tau^*) - \pi_p^*(a^*)] dx$$

as well as the dual functional

$$J^*[a^*, \tau^*] := \inf_{w \in \mathbb{K}} \ell(w, a^*, \tau^*),$$

then the dual problem

$$J^* \rightarrow \max \text{ on } X^* \times Y^*$$

has a unique solution (d^*, σ^*) for which

$$(2.1) \quad J[u] = J^*[d^*, \sigma^*],$$

u denoting the solution of (\mathcal{P}) , we refer again to [ET]. As done in [Re] we define suitable perturbations of problem (\mathcal{P}) : for $\lambda \in \Lambda := \{\rho \in L^q(\Omega) : \rho \geq 0\}$ we let

$$(\mathcal{P}_{\lambda}) \quad J_{\lambda}[w] := J[w] - \int_{\Omega} \lambda(w - \Psi) dx \rightarrow \min \text{ in } W_p^2(\Omega) \cap \overset{\circ}{W}_p^1(\Omega)$$

and observe that (\mathcal{P}_{λ}) admits a unique solution u_{λ} . Moreover, it is immediate that

$$\begin{aligned} \sup_{\lambda \in \Lambda} J_{\lambda}[w] &= J[w] - \inf_{\lambda \in \Lambda} \int_{\Omega} \lambda(w - \Psi) dx \\ &= \begin{cases} J[w], & \text{if } w \in \mathbb{K}, \\ +\infty, & \text{if } w \notin \mathbb{K}. \end{cases} \end{aligned}$$

The Lagrangian associated to J_{λ} is given by

$$\begin{aligned} L(w, a^*, \tau^*, \lambda) &= \ell(w, a^*, \tau^*) - \int_{\Omega} \lambda(w - \Psi) dx, \\ w &\in W_p^2(\Omega) \cap \overset{\circ}{W}_p^1(\Omega), (a^*, \tau^*) \in X^* \times Y^*, \lambda \in \Lambda, \end{aligned}$$

and the maximizing problem

$$J_\lambda^*[a^*, \xi^*] := \inf_{w \in \mathring{W}_p^1(\Omega) \cap W_p^2(\Omega)} L(w, a^*, \xi^*, \lambda) \rightarrow \max \text{ in } X^* \times Y^*$$

dual to (\mathcal{P}_λ) has a unique solution $(d_\lambda^*, \sigma_\lambda^*)$ such that

$$(2.2) \quad J_\lambda[u_\lambda] = J_\lambda^*[d_\lambda^*, \sigma_\lambda^*].$$

We remark that $J_\lambda^*[a^*, \tau^*] > -\infty$ implies that

$$(2.3) \quad \int_\Omega [a^* \cdot \nabla w + \tau^* : \nabla^2 w - \lambda w] dx = 0$$

for all $w \in W_p^2(\Omega) \cap \mathring{W}_p^1(\Omega)$. Let $Q_\lambda^* := \{(a^*, \tau^*) \in X^* \times Y^* : (a^*, \tau^*) \text{ satisfies (2.3)}\}$.

Taking $w \in \mathring{W}_p^2(\Omega)$ in (2.3) we see that in the distributional sense

$$\operatorname{div}(\operatorname{div} \tau^*) - \operatorname{div} a^* = \lambda$$

is true for $(a^*, \tau^*) \in Q_\lambda^*$. We further observe the following inequality:

$$\inf_{v \in \mathring{W}_p^1(\Omega) \cap W_p^2(\Omega)} J_\lambda[v] \leq \inf_{w \in \mathbb{K}} J_\lambda[w] = \inf_{w \in \mathbb{K}} [J[w] - \int_\Omega \lambda(w - \Psi) dx] \leq \inf_{w \in \mathbb{K}} J[w],$$

or equivalently:

$$(2.4) \quad J_\lambda[u_\lambda] \leq J[u].$$

After these preparations we can state our first result:

THEOREM 2.1. *Let $p \geq 2$. With the notation introduced above we have for any $v \in \mathbb{K}$, for any $b^* \in X^*$, for all $\xi^* \in Y^*$, for any $\lambda \in \Lambda$ and for any choice of $\beta > 0$ the estimate*

$$(2.5) \quad \begin{aligned} & \|\nabla^2(u - v)\|_{L^p}^p + \|\nabla(u - v)\|_{L^p}^p \\ & \leq p \cdot 2^{p-1} \{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + [2^{2-q}(3-q) + \frac{1}{q}\beta^{-q}] d(b^*, \xi^*)^q \\ & \quad + \frac{1}{p}\beta^p [\|\xi^*\|^{q-2} \xi^* - \nabla^2 v \|_{L^p}^p + \| |b^*|^{q-2} b^* - \nabla v \|_{L^p}^p] \\ & \quad + \int_\Omega \lambda(v - \Psi) dx \}. \end{aligned}$$

Here we have abbreviated $D_p : X \times Y \times X^* \times Y^* \rightarrow [0, \infty)$,

$$\begin{aligned} D_p[a, \eta, a^*, \eta^*] & := \int_\Omega [\pi_p(a) + \pi_p^*(a^*) - a^* \cdot a] dx + \int_\Omega [\Pi_p(\eta) + \Pi_p^*(\eta^*) - \eta : \eta^*] dx, \\ d(b^*, \eta^*)^q & := \inf_{(a^*, \tau^*) \in Q_\lambda^*} [\|\tau^* - \eta^*\|_{L^q}^q + \|a^* - b^*\|_{L^q}^q]. \end{aligned}$$

REMARK 2.1. *i) An estimate for the distance $d(b^*, \eta^*)$ of (b^*, η^*) to Q_λ^* can be obtained along the lines of [BFR2], Theorem 3.1 and Theorem 3.2.*

ii) Following [Re] we can choose the function λ in natural ways in order to get variants of Corollary 3.1 and 3.2 from [BFR2].

iii) If $p < 2$, then one can follow Section 4 of [BFR2] to find the appropriate version of Theorem 2.1 in which the minimizer u is replaced by the maximizer (d^, σ^*) .*

iv) Clearly all the terms on the r.h.s. of (2.5) are non-negative, and they vanish simultaneously if and only if $(b^, \xi^*) \in Q_\lambda^*$, $\lambda(v - \Psi) = 0$, $\nabla^2 v = |\xi^*|^{q-2} \xi^*$, $\nabla v = |b^*|^{q-2} b^*$.*

Proof of Theorem 2.1: Let $v \in \mathbb{K}$, $\lambda \in \Lambda$ and $(a^*, \tau^*) \in Q_\lambda^*$. For vector- or tensor-valued functions A, B of class $L^p(\Omega)$ we have by the variant of Clarkson's inequality [Cl] proved in [MM] on account of $p \geq 2$

$$(2.6) \quad \int_{\Omega} \left[\left| \frac{A+B}{2} \right|^p + \left| \frac{A-B}{2} \right|^p \right] dx \leq \frac{1}{2} \|A\|_{L^p}^p + \frac{1}{2} \|B\|_{L^p}^p.$$

Applying (2.6) in an obvious way we see

$$\|\nabla^2(u-v)\|_{L^p}^p + \|\nabla(u-v)\|_{L^p}^p \leq p \, 2^p \left[\frac{1}{2} J[v] + \frac{1}{2} J[u] - J\left[\frac{u+v}{2}\right] \right],$$

and the minimality of u implies

$$(2.7) \quad \|\nabla^2(u-v)\|_{L^p}^p + \|\nabla(u-v)\|_{L^p}^p \leq p \, 2^{p-1} \left[J[v] - J[u] \right].$$

From (2.2) and (2.4) we get

$$J[u] \geq J_\lambda^*[d_\lambda^*, \sigma_\lambda^*] \geq J_\lambda^*[a^*, \tau^*]$$

which gives in combination with (2.7)

$$(2.8) \quad \|\nabla^2(u-v)\|_{L^p}^p + \|\nabla(u-v)\|_{L^p}^p \leq p \, 2^{p-1} \left[J[v] - J_\lambda^*[a^*, \tau^*] \right].$$

We discuss the r.h.s. of (2.8): for $(a^*, \tau^*) \in Q_\lambda^*$ we have

$$J_\lambda^*[a^*, \tau^*] = \int_{\Omega} \left[-\Pi_p^*(\tau^*) - \pi_p^*(a^*) + \Psi \lambda \right] dx,$$

hence

$$\begin{aligned} J[v] - J_\lambda^*[a^*, \tau^*] &= \int_{\Omega} \left[\Pi_p(\nabla^2 v) + \Pi_p^*(\tau^*) - \tau^* : \nabla^2 v \right. \\ &\quad \left. + \pi_p(\nabla v) + \pi_p^*(a^*) - a^* \cdot \nabla v \right] dx + \int_{\Omega} (v - \Psi) \lambda dx \\ &= D_p[\nabla v, \nabla^2 v, a^*, \tau^*] + \int_{\Omega} (v - \Psi) \lambda dx. \end{aligned}$$

Inserting this into (2.8) it is shown that

$$(2.9) \quad \|\nabla^2(u-v)\|_{L^p}^p + \|\nabla(u-v)\|_{L^p}^p \leq p 2^{p-1} \left\{ D_p[\nabla v, \nabla^2 v, a^*, \tau^*] + \int_{\Omega} (v - \Psi)\lambda dx \right\}$$

for all $v \in \mathbb{K}$, $(a^*, \tau^*) \in Q_{\lambda}^*$, $\lambda \in \Lambda$. Next we let $b^* \in X^*$, $\xi^* \in Y^*$. If v, a^*, τ^*, λ are as in (2.9), we get by the convexity of Π_p^* , π_p^* and by(2.9)

$$\begin{aligned} & \|\nabla^2(u-v)\|_{L^p}^p + \|\nabla(u-v)\|_{L^p}^p \\ & \leq p 2^{p-1} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + \int_{\Omega} [\Pi_p^*(\tau^*) - \Pi_p^*(\xi^*) - (\tau^* - \xi^*) : \nabla^2 v] dx \right. \\ & \quad \left. + \int_{\Omega} [\pi_p^*(a^*) - \pi_p^*(b^*) - (a^* - b^*) \cdot \nabla v] dx + \int_{\Omega} (v - \Psi)\lambda dx \right\} \\ & \leq p 2^{p-1} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + \int_{\Omega} [|\tau^*|^{q-2}\tau^* - \nabla^2 v] : (\tau^* - \xi^*) dx \right. \\ & \quad \left. + \int_{\Omega} [|a^*|^{q-2}a^* - \nabla v] \cdot (a^* - b^*) dx + \int_{\Omega} (v - \Psi)\lambda dx \right\} \\ & = p 2^{p-1} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + \int_{\Omega} [|\tau^*|^{q-2}\tau^* - |\xi^*|^{q-2}\xi^*] : (\tau^* - \xi^*) dx \right. \\ & \quad \left. + \int_{\Omega} [|a^*|^{q-2}a^* - |b^*|^{q-2}b^*] \cdot (a^* - b^*) dx + \int_{\Omega} [|\xi^*|^{q-2}\xi^* - \nabla^2 v] : (\tau^* - \xi^*) dx \right. \\ & \quad \left. + \int_{\Omega} [|b^*|^{q-2}b^* - \nabla v] \cdot (a^* - b^*) dx + \int_{\Omega} \lambda(v - \Psi) dx \right\} \\ & =: p 2^{p-1} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + \sum_{i=1}^4 I_i + \int_{\Omega} \lambda(v - \Psi) dx \right\}. \end{aligned}$$

According to [BR] we have

$$\begin{aligned} I_1 & \leq 2^{2-q}(3-q)\|\tau^* - \xi^*\|_{L^q}^q, \\ I_2 & \leq 2^{2-q}(3-q)\|a^* - b^*\|_{L^q}^q, \end{aligned}$$

and from Hölder's and Young's inequality we find

$$\begin{aligned} I_3 & \leq \| |\xi^*|^{q-2}\xi^* - \nabla^2 v \|_{L^p} \|\tau^* - \xi^*\|_{L^q} \\ & \leq \frac{1}{p} \beta^p \| |\xi^*|^{q-2}\xi^* - \nabla^2 v \|_{L^p}^p + \frac{1}{q} \beta^{-q} \|\tau^* - \xi^*\|_{L^q}^q, \\ I_4 & \leq \frac{1}{p} \beta^p \| |b^*|^{q-2}b^* - \nabla v \|_{L^p}^p + \frac{1}{q} \beta^{-q} \|a^* - b^*\|_{L^q}^q, \end{aligned}$$

where $\beta > 0$ is arbitrary. Collecting the various estimates it is shown that

$$\begin{aligned} & \|\nabla^2(v-u)\|_{L^p}^p + \|\nabla(v-u)\|_{L^p}^p \\ & \leq p 2^{p-1} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + 2^{2-q}(3-q) [\|\tau^* - \xi^*\|_{L^q}^q + \|a^* - b^*\|_{L^q}^q] \right. \\ & \quad + \frac{1}{p} \beta^p [\|\xi^*|^{q-2}\xi^* - \nabla^2 v\|_{L^p}^p + \| |b^*|^{q-2}b^* - \nabla v \|_{L^p}^p] \\ & \quad \left. + \frac{1}{q} \beta^{-q} [\|\tau^* - \xi^*\|_{L^q}^q + \|a^* - b^*\|_{L^q}^q] + \int_{\Omega} \lambda(v - \Psi) dx \right\}, \end{aligned}$$

and inequality (2.5) follows by taking the inf w.r.t. $(a^*, \tau^*) \in Q_{\lambda}^*$. \square

3 An estimate for the distance to the set of admissible comparison functions

In order to apply inequality (2.5) we have to take functions v from the class of admissible comparison functions. Now, if $w \in \overset{\circ}{W}_p^1(\Omega) \cap W_p^2(\Omega)$ is arbitrary we can not simply “project” w on the class \mathbb{K} as it is possible for first order variational inequalities since $\max(w, \Psi)$ in general is not an element of $W_p^2(\Omega)$. So it remains to measure the distance of w to the set \mathbb{K} , and a reasonable quantity to do this is given by

$$\rho(w)^p := \inf_{v' \in \mathbb{K}} \left[\int_{\Omega} \Pi_p(\nabla^2 v' - \nabla^2 w) + \pi_p(\nabla v' - \nabla w) \right] dx.$$

If $v' \in \mathbb{K}$, then $v' - w \in W_p^2(\Omega) \cap \overset{\circ}{W}_p^1(\Omega)$ together with $v' - w \geq \tilde{\Psi} := \Psi - w$, hence

$$(3.1) \quad \rho(w)^p = \inf \{ J[v] : v \in W_p^2(\Omega) \cap \overset{\circ}{W}_p^1(\Omega), v \geq \tilde{\Psi} \}.$$

As a comparison function for the minimization problem on the r.h.s. of (3.1) we consider the solution h of the first order variational inequality

$$\int_{\Omega} |\nabla g|^2 dx \rightarrow \min \text{ in } \{ f \in \overset{\circ}{W}_2^1(\Omega) : f \geq \tilde{\Psi} \}.$$

h solves the equation

$$-\Delta h = \begin{cases} 0 & \text{on } [h > \tilde{\Psi}] \\ -\Delta \tilde{\Psi} & \text{on } [h = \tilde{\Psi}] \end{cases} =: F \in L^p(\Omega),$$

and by the L^p -theory for elliptic equations (see, e.g. [Mo], Theorem 5.6.2, or [GT]) h is in the space $W_p^2(\Omega) \cap \overset{\circ}{W}_p^1(\Omega)$ together with

$$(3.2) \quad \int_{\Omega} |\nabla^2 h|^p dx \leq C_1(p, \Omega) \int_{\Omega} |F|^p dx.$$

On the other hand, the interpolation theorem 4.14 of [Ad] shows

$$(3.3) \quad \int_{\Omega} |\nabla h|^p dx \leq C_2(p, \Omega) \left[\int_{\Omega} |\nabla^2 h|^p dx + \int_{\Omega} |h|^p dx \right].$$

For discussing $\int_{\Omega} |h|^p dx$ we assume $p > 2$ and let $r := 2p/(p+2)$. Since $h \in \mathring{W}_2^1(\Omega)$ and $r < 2$, we can apply Sobolev's and Hölder's inequality to get

$$\begin{aligned} \int_{\Omega} |h|^p dx &\leq C_3(p, \Omega) \left(\int_{\Omega} |\nabla h|^r dx \right)^{p/r} \leq C_4(p, \Omega) \left(\int_{\Omega} |\nabla h|^2 dx \right)^{p/2} \\ &= C_4(p, \Omega) \left(\int_{\Omega} hF dx \right)^{p/2} = C_4(p, \Omega) \left(\int_{[h=\tilde{\Psi}]} |\tilde{\Psi}\Delta\tilde{\Psi}| dx \right)^{p/2}. \end{aligned}$$

Combining this estimate with (3.2) and (3.3) it is shown that

$$(3.4) \quad \begin{aligned} \rho(w)^p &\leq C_5(p, \Omega) \left[\int_{\Omega} |F|^p dx + \left(\int_{[h=\tilde{\Psi}]} |\tilde{\Psi}\Delta\tilde{\Psi}| dx \right)^{p/2} \right] \\ &= C_5(p, \Omega) \left[\int_{[h=\tilde{\Psi}]} |\Delta\tilde{\Psi}|^p dx + \left(\int_{[h=\tilde{\Psi}]} |\tilde{\Psi}\Delta\tilde{\Psi}| dx \right)^{p/2} \right] \end{aligned}$$

In order to proceed further we observe that $-\Delta h \geq 0$ which is an immediate consequence of

$$\int_{\Omega} |\nabla h|^2 dx \leq \int_{\Omega} |\nabla h + t\nabla\eta|^2 dx$$

valid for $t \geq 0$ and all $\eta \in C_0^1(\Omega)$, $\eta \geq 0$. Since $h|_{\partial\Omega} = 0$ we deduce $h \geq 0$ on Ω , hence $[h = \tilde{\Psi}] \subset [\Psi - w \geq 0] = [w \leq \Psi]$. Inserting this into (3.4) we have shown:

THEOREM 3.1. *If $w \in W_p^2(\Omega) \cap \mathring{W}_p^1(\Omega)$ is arbitrary, then in case $p \geq 2$ we have with a suitable constant $C = C(p, \Omega) > 0$*

$$(3.5) \quad \begin{aligned} &\inf_{v' \in \mathbb{K}} \left[\|\nabla^2 w - \nabla^2 v'\|_{L^p}^p + \|\nabla w - \nabla v'\|_{L^p}^p \right] \\ &\leq C \left[\int_{[w \leq \Psi]} |\Delta(\Psi - w)|^p dx + \left(\int_{[w \leq \Psi]} |\Psi - w| |\Delta(\Psi - w)| dx \right)^{p/2} \right]. \end{aligned}$$

REMARK 3.1. *A similar estimate is valid in case $p < 2$.*

Theorem 3.1 is applied as follows: let $w \in W_p^2(\Omega) \cap \mathring{W}_p^1(\Omega)$ and consider $b^* \in X^*$, $\xi^* \in Y^*$, $\lambda \in \Lambda$ and $\beta > 0$. Then (2.5) gives for any $v \in \mathbb{K}$

$$\begin{aligned} &\|\nabla^2(u - w)\|_{L^p}^p + \|\nabla(u - w)\|_{L^p}^p \\ &\leq 2^{p-1} [\|\nabla^2(u - v)\|_{L^p}^p + \|\nabla(u - v)\|_{L^p}^p] + 2^{p-1} [\|\nabla^2(w - v)\|_{L^p}^p + \|\nabla(w - v)\|_{L^p}^p] \\ &\leq p 2^{2p-2} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] + [2^{2-q}(3-q) + \frac{1}{q} \beta^{-q}] d(b^*, \xi^*)^q \right. \\ &\quad \left. + \frac{1}{p} \beta^p [|\xi^*|^{q-2} \xi^* - \nabla^2 v]_{L^p}^p + [|b^*|^{q-2} b^* - \nabla v]_{L^p}^p + \int_{\Omega} \lambda(v - \Psi) dx \right\} \\ &\quad + 2^{p-1} [\|\nabla^2(w - v)\|_{L^p}^p + \|\nabla(w - v)\|_{L^p}^p]. \end{aligned}$$

If we replace the function v in $\{\dots\}$ by the function w and estimate the resulting difference in an obvious way, then we arrive at

$$\begin{aligned}
& \|\nabla^2(u-w)\|_{L^p}^p + \|\nabla(u-w)\|_{L^p}^p \\
& \leq p 2^{2p-2} \left\{ D_p[\nabla w, \nabla^2 w, b^*, \xi^*] + [2^{2-q}(3-q) + \frac{1}{q} \beta^{-q}] d(b^*, \xi^*)^q \right. \\
& \quad + \frac{2^{p-1}}{p} \beta^p [\|\xi^*|^{q-2} \xi^* - \nabla^2 w\|_{L^p}^p + \||b^*|^{q-2} b^* - \nabla w\|_{L^p}^p] \\
& \quad \left. + \int_{\Omega} \lambda(w - \Psi) dx \right\} + p 2^{2p-2} \|\lambda\|_{L^q} \|w - v\|_{L^p} \\
& \quad + (2^{p-1} + \beta^p 2^{3p-3}) [\|\nabla^2(w-v)\|_{L^p}^p + \|\nabla(w-v)\|_{L^p}^p] \\
& \quad + p 2^{2p-2} \left\{ D_p[\nabla v, \nabla^2 v, b^*, \xi^*] - D_p[\nabla w, \nabla^2 w, b^*, \xi^*] \right\}.
\end{aligned}$$

We observe

$$\begin{aligned}
& D_p[\nabla v, \nabla^2 v, b^*, \xi^*] - D_p[\nabla w, \nabla^2 w, b^*, \xi^*] \\
& = \int_{\Omega} [\pi_p(\nabla v) - \pi_p(\nabla w) - b^* \cdot (\nabla v - \nabla w)] dx \\
& \quad + \int_{\Omega} [\Pi_p(\nabla^2 v) - \Pi_p(\nabla^2 w) - \xi^* : (\nabla^2 v - \nabla^2 w)] dx \\
& \leq \int_{\Omega} (D\pi_p(\nabla v) - b^*) \cdot (\nabla v - \nabla w) dx \\
& \quad + \int_{\Omega} (D\Pi_p(\nabla^2 v) - \xi^*) : (\nabla^2 v - \nabla^2 w) dx \\
& = \int_{\Omega} (D\pi_p(\nabla v) - D\pi_p(\nabla w)) \cdot (\nabla v - \nabla w) dx \\
& \quad + \int_{\Omega} (D\pi_p(\nabla w) - b^*) \cdot (\nabla v - \nabla w) dx \\
& \quad + \int_{\Omega} (D\Pi_p(\nabla^2 v) - D\Pi_p(\nabla^2 w)) : (\nabla^2 v - \nabla^2 w) dx \\
& \quad + \int_{\Omega} (D\Pi_p(\nabla^2 w) - \xi^*) : (\nabla^2 v - \nabla^2 w) dx
\end{aligned}$$

where we used the convexity of the potentials. We have

$$\begin{aligned}
& \int_{\Omega} (D\pi_p(\nabla w) - b^*) \cdot (\nabla v - \nabla w) dx + \int_{\Omega} (D\Pi_p(\nabla^2 w) - \xi^*) : (\nabla^2 v - \nabla^2 w) dx \\
& \leq \|\nabla w\|_{L^q}^{p-2} \|\nabla w - b^*\|_{L^q} \|\nabla v - \nabla w\|_{L^p} + \|\nabla^2 w\|_{L^q} \|\nabla^2 w - \xi^*\|_{L^q} \|\nabla^2 v - \nabla^2 w\|_{L^p}
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\Omega} (D\pi_p(\nabla w) - D\pi_p(\nabla v)) \cdot (\nabla v - \nabla w) dx \\
& + \int_{\Omega} (D\Pi_p(\nabla^2 w) - D\Pi_p(\nabla^2 v)) : (\nabla^2 v - \nabla^2 w) dx \\
& \leq \{ \|D\pi_p(\nabla v)\|_{L^q} + \|D\pi_p(\nabla w)\|_{L^q} \} \|\nabla v - \nabla w\|_{L^p} \\
& + \{ \|D\Pi_p(\nabla^2 v)\|_{L^q} + \|D\Pi_p(\nabla^2 w)\|_{L^q} \} \|\nabla^2 v - \nabla^2 w\|_{L^p} \\
& = \{ \|\nabla v\|_{L^p}^{p-1} + \|\nabla w\|_{L^p}^{p-1} \} \|\nabla v - \nabla w\|_{L^p} + \{ \|\nabla^2 v\|_{L^p}^{p-1} + \|\nabla^2 w\|_{L^p}^{p-1} \} \|\nabla^2 v - \nabla^2 w\|_{L^p} \\
& \leq [2^{p-2} \|\nabla v - \nabla w\|_{L^p}^{p-1} + 2^{p-2} \|\nabla w\|_{L^p}^{p-1} + \|\nabla w\|_{L^p}^{p-1}] \|\nabla v - \nabla w\|_{L^p} \\
& + [2^{p-2} \|\nabla^2 v - \nabla^2 w\|_{L^p}^{p-1} + 2^{p-2} \|\nabla^2 w\|_{L^p}^{p-1} + \|\nabla^2 w\|_{L^p}^{p-1}] \|\nabla^2 v - \nabla^2 w\|_{L^p} \\
& = 2^{p-2} [\|\nabla v - \nabla w\|_{L^p}^p + \|\nabla^2 v - \nabla^2 w\|_{L^p}^p] \\
& + (2^{p-2} + 1) [\|\nabla w\|_{L^p}^{p-1} \|\nabla v - \nabla w\|_{L^p} + \|\nabla^2 w\|_{L^p}^{p-1} \|\nabla^2 v - \nabla^2 w\|_{L^p}].
\end{aligned}$$

Collecting terms we get (recalling Theorem 3.1)

THEOREM 3.2. *For any $w \in W_p^2(\Omega) \cap \dot{W}_p^1(\Omega)$, $b^* \in X^*$, $\xi^* \in Y^*$, $\lambda \in \Lambda$ and $\beta > 0$ it holds:*

$$\begin{aligned}
& \|\nabla^2(u - w)\|_{L^p}^p + \|\nabla(u - w)\|_{L^p}^p \\
& \leq 2^{2p-2} [r.h.s. \text{ of (2.5) with } v \text{ replaced by } w] + p 2^{2p-2} \|\lambda\|_{L^q} \|w - v\|_{L^p} \\
& + [2^{p-1} + \beta^p 2^{3p-3} + p 2^{3p-4}] [\|\nabla^2 v - \nabla^2 w\|_{L^p}^p + \|\nabla v - \nabla w\|_{L^p}^p] \\
& + p 2^{2p-2} [\|\nabla w\|_{L^q}^{p-2} \|\nabla w - b^*\|_{L^q} \|\nabla v - \nabla w\|_{L^p} \\
& + \|\nabla^2 w\|_{L^q}^{p-2} \|\nabla^2 w - \xi^*\|_{L^q} \|\nabla^2 v - \nabla^2 w\|_{L^p}] \\
(3.6) \quad & + p 2^{2p-2} (2^{p-2} + 1) [\|\nabla w\|_{L^p}^{p-1} \|\nabla v - \nabla w\|_{L^p} + \|\nabla^2 w\|_{L^p}^{p-1} \|\nabla^2 v - \nabla^2 w\|_{L^p}].
\end{aligned}$$

Here v denotes any function from the class \mathbb{K} , and in the above inequality we may replace $\|\nabla^i v - \nabla^i w\|_{L^p}$, $i = 0, 1, 2$, by $\mathcal{R}(w, \Psi)^{1/p}$, $\mathcal{R}(w, \Psi)$ denoting the r.h.s. of the inequality (3.5).

REMARK 3.2. *i) Since we use the Poincaré-inequality for the term $\|w - v\|_{L^p}$, the constant C appearing in (3.5) has to be adjusted to ensure the last statement of Theorem 3.2.*

ii) Note that after taking the inf w.r.t. $v \in \mathbb{K}$ inequality (3.6) reduces to (2.5) with a slightly larger factor in front of $\{\dots\}$ on the r.h.s. provided we start from a function $w \in \mathbb{K}$. Inequality (2.5) is not exactly reproduced since the expression

$$\|\nabla \dots\|_{L^p}^p + \|\nabla^2 \dots\|_{L^p}^p$$

is not a norm. Replacing it by

$$\|\nabla \dots\|_{L^p} + \|\nabla^2 \dots\|_{L^p}$$

would cause the same difficulty in a different place of the calculations.

References

- [Ad] Adams, R. A., Sobolev spaces. Academic Press, New York-San Francisco-London 1975.
- [BFR1] Bildhauer, M., Fuchs, M., Repin, S., A posteriori error estimates for stationary slow flows of power-law fluids. *J. Non-Newtonian Fluid Mech.* 142 (2007), 112–122.
- [BFR2] Bildhauer, M., Fuchs, M., Repin, S., Duality based a posteriori error estimates for higher order variational inequalities with power growth functionals. Submitted.
- [BR] Bildhauer, M., Repin, S., Estimates for the deviation from exact solutions of variational problems with power growth functionals. *Zap. Nauch. Semi. (POMI)* 336 (2006), 5–24.
- [Cl] Clarkson, J.A., Uniformly convex spaces. *Trans. Am. Math. Soc.* 40 (1936), 396–414.
- [ET] Ekeland, I., Temam, R., Convex analysis and variational problems. North-Holland, Amsterdam, 1976.
- [GT] Gilbarg, D., Trudinger, N.S., Elliptic partial differential equations of second order. *Grundlehren der math. Wiss.* 224, second ed., revised third print., Springer, Berlin-Heidelberg-New York 1998.
- [Mo] Morrey, C. B., Multiple integrals in the calculus of variations. *Grundlehren der math. Wiss. in Einzeldarstellungen* 130, Springer, Berlin–Heidelberg–New York 1966.
- [MM] P.P. Mosolov and V.P. Mjasnikov, Mechanics of rigid plastic media. Nauka, Moscow (1981) (in Russian).
- [NR] Neittaanmäki, P., Repin, S., Reliable methods for computer simulation: error control and a posteriori estimates. Elsevier, New York, 2004.
- [Re] Repin, S., Estimates of deviations from exact solutions of elliptic variational inequalities. *J. Math. Sci.* 115 (2003), 2811–2819.